



Test Definition: FFMI2

Anti-Mi-2 Ab

Overview

Method Name

RIPA Gel Radiography

NY State Available

Yes

Specimen

Specimen Type

Serum

Specimen Required

Collection Container/Tube:

Preferred: Red top

Acceptable: Serum gel

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL serum

Collection Instructions:

1. Within 1 hour of collection, centrifuge and aliquot 1 mL of serum into a plastic vial.
2. Send refrigerate.

Specimen Minimum Volume

Serum: 0.5 mL Note: This volume does not allow for repeat testing.

Reject Due To

Gross hemolysis	Reject
Gross lipemia	Reject
Gross icterus	Reject
Specimens other than serum	Reject
Bacterial contamination	Reject

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
---------------	-------------	------	-------------------

Serum	Refrigerated (preferred)	14 days	
	Ambient	7 days	
	Frozen	60 days	

Clinical & Interpretive

Clinical Information

Refer to www.labcorp.com/test-menu

Reference Values

Negative

Performance

PDF Report

No

Day(s) Performed

Daily

Report Available

16 to 25 days

Performing Laboratory Location

Esoterix Endocrinology

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics validated by LabCorp. It has not been cleared or approved by the Food and Drug Administration.

CPT Code Information

83516

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
FFMI2	Anti-Mi-2 Ab	18485-3

Result ID	Test Result Name	Result LOINC® Value
FFMI2	Anti-Mi-2 Ab	18485-3