MAYO CLINIC LABORATORIES

Test Definition: BUN

Blood Urea Nitrogen (BUN), Serum

Overview

Useful For
Screening test for evaluation of kidney function

Method Name
Photometric

NY State Available
Yes

Specimen

Specimen Type
Serum

Necessary Information
Patient's age and sex are required.

Specimen Required
Collection Container/Tube:
Preferred: Serum gel
Acceptable: Red top
Submission Container/Tube: Plastic vial
Specimen Volume: 0.5 mL
Collection Instructions:
1. Serum gel tubes should be centrifuged within 2 hours of collection.
2. Red-top tubes should be centrifuged, and the serum aliquoted into a plastic vial within 2 hours of collection.

Forms
If not ordering electronically, complete, print, and send 1 of the following forms with the specimen:
-Kidney Transplant Test Request
-Renal Diagnostics Test Request (T830)

Specimen Minimum Volume
See Specimen Required

Reject Due To

Gross hemolysis	OK

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	7 days	
	Frozen	365 days	

Clinical & Interpretive

Clinical Information

Urea is the final degradation product of protein and amino acid metabolism. In protein catabolism, the proteins are broken down to amino acids and deaminated. The ammonia formed in this process is synthesized to urea in the liver. This is the most important catabolic pathway for eliminating excess nitrogen in the human body.

Increased blood urea nitrogen (BUN) may be due to prerenal causes (cardiac decompensation, water depletion due to decreased intake and excessive loss, increased protein catabolism, and high protein diet), renal causes (acute glomerulonephritis, chronic nephritis, polycystic kidney disease, nephrosclerosis, and tubular necrosis), and postrenal causes (eg, all types of obstruction of the urinary tract, such as stones, enlarged prostate gland, tumors).

The determination of serum BUN currently is the most widely used screening test for the evaluation of kidney function. The test is frequently requested along with the serum creatinine test since simultaneous determination of these 2 compounds appears to aid in the differential diagnosis of prerenal, renal and postrenal hyperuremia.

Reference Values

Males
1-17 years: 7-20 mg/dL
> or =18 years: 8-24 mg/dL
Reference values have not been established for patients who are <12 months of age.

Females
1-17 years: 7-20 mg/dL
> or =18 years: 6-21 mg/dL
Reference values have not been established for patients who are <12 months of age.

Cautions

No significant cautionary statements.

Clinical Reference

Lamb EJ, Jones GRD: Kidney function tests. In: Rifai N, Horvath AR, Wittwer CT, eds. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics. 6th ed. Elsevier; 2018:497-500

Performance

Method Description

Kinetic test with urease and glutamate dehydrogenase. Urea is hydrolyzed by urease to form ammonium and carbonate. In the second reaction 2-oxoglutarate reacts with ammonium in the presence of glutamate dehydrogenase and the coenzyme reduced nicotinamide adenine dinucleotide (NADH) to produce L-glutamate. In this reaction two moles of NADH are oxidized to NAD[+] for each mole of urea hydrolyzed. The rate of decrease in NADH concentrations is directly proportional to the urea concentration in the specimen and is measured photometrically.(Package insert: Urea/BUN reagent. Roche Diagnostics; 07/2024)

PDF Report
No

Day(s) Performed
Monday through Sunday

Report Available
Same day/1 to 2 days

Specimen Retention Time
1 week

Performing Laboratory Location
Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees
- Authorized users can sign in to Test Prices for detailed fee information.
- Clients without access to Test Prices can contact Customer Service 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact Customer Service.

Test Classification
This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information
84520

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
BUN	Bld Urea Nitrog(BUN), S	3094-0

Result ID	Test Result Name	Result LOINC® Value
BUN	Bld Urea Nitrog(BUN), S	3094-0