Test Definition: MOLD1

Mold Panel, Serum

Overview

Useful For

Establishing a diagnosis of an allergy to mold

Defining the allergen responsible for eliciting signs and symptoms

Identifying allergens:
-Responsible for allergic response and/or anaphylactic episode
-To confirm sensitization prior to beginning immunotherapy
-To investigate the specificity of allergic reactions to insect venom allergens, drugs, or chemical allergens

Testing for IgE antibodies is **not useful** in patients previously treated with immunotherapy to determine if residual clinical sensitivity exists or in patients in whom the medical management does not depend upon identification of allergen specificity.

Testing Algorithm

Includes testing for *Alternaria tenuis*, *Aspergillus fumigatus*, *Candida albicans* (monilia), *Cladosporium herbarum*, *Helminthosporium halodes*, and *Penicillium chrysogenum* allergen.

Special Instructions

- Allergens - Immunoglobulin E (IgE) Antibodies

Highlights

This multi-allergen IgE antibody panel, combined with measurement of IgE in serum, is an appropriate first-order test for allergic disease.

It requires less specimen volume and less cost for ruling out allergic response; however, individual (single) allergen responses cannot be identified. In cases of a positive test, follow-up testing must be performed to differentiate between individual allergens in the panel.

Note: Only one result is generated for each panel.

Method Name

Fluorescence Enzyme Immunoassay (FEIA)

NY State Available

Yes

Specimen

Specimen Type

Serum

Ordering Guidance

This test uses a pooled allergen reagent; therefore, the multi-allergen Immunocap (panel cap) is reported with a single qualitative class result and concentration. This is the appropriate first-tier test for allergic disease.

For a listing of allergens available for testing, see Allergens - Immunoglobulin E (IgE) Antibodies.

Specimen Required

Collection Container/Tube:
Preferred: Serum gel
Acceptable: Red top
Submission Container/Tube: Plastic vial
Specimen Volume: 0.5 mL for every 5 allergens requested
Collection Instructions: Centrifuge and aliquot serum into a plastic vial.

Forms

If not ordering electronically, complete, print, and send an Allergen Test Request (T236) with the specimen.

Specimen Minimum Volume

For 1 allergen: 0.3 mL
For more than 1 allergen: (0.05 mL x number of allergens) + 0.25 mL deadspace

Reject Due To

Gross hemolysis	OK
Gross lipemia	OK

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	14 days	
	Frozen	90 days	

Clinical & Interpretive

Clinical Information

Clinical manifestations of immediate hypersensitivity (allergic) diseases are caused by the release of proinflammatory mediators (histamine, leukotrienes, and prostaglandins) from IgE-sensitized effector cells (mast cells and basophils) when cell-bound IgE antibodies interact with an allergen.

In vitro serum testing for IgE antibodies provides an indication of the immune response to allergens that may be

associated with allergic disease.

The allergens chosen for testing often depend upon the age of the patient, history of allergen exposure, season of the year, and clinical manifestations. In individuals predisposed to develop allergic disease, the sequence of sensitization and clinical manifestations proceed as follows: eczema and respiratory disease (rhinitis and bronchospasm) in infants and children younger than 5 years due to food sensitivity (milk, egg, soy, and wheat proteins) followed by respiratory disease (rhinitis and asthma) in older children and adults due to sensitivity to inhalant allergens (dust mite, mold, and pollen inhalants).

Reference Values

Class	IgE kU/L	Interpretation
0	<0.10	Negative
0/1	0.10-0.34	Borderline/equivocal
1	0.35-0.69	Equivocal
2	0.70-3.49	Positive
3	3.50-17.4	Positive
4	17.5-49.9	Strongly positive
5	50.0-99.9	Strongly positive
6	> or =100	Strongly positive

Reference values apply to all ages.

Concentrations of 0.70 kU/L or more (class 2 and above) will flag as abnormally high.

Interpretation

Positive results indicate the possibility of allergic disease induced by one or more allergens present in the multi-allergen cap.

Negative results may rule out allergy, except in rare cases of allergic disease induced by exposure to a single allergen.

Detection of IgE antibodies in serum (class 1 or greater) indicates an increased likelihood of allergic disease as opposed to other etiologies and defines the allergens that may be responsible for eliciting signs and symptoms.

The level of IgE antibodies in serum varies directly with the concentration of IgE antibodies expressed as a class score or kU/L.

Cautions

Positive results indicate the possibility of allergic disease induced by one or more allergens present in the multi-allergen cap.

Negative results may rule out allergy, except in rare cases of allergic disease induced by exposure to a single allergen.

Some individuals with clinically insignificant sensitivity to allergens may have measurable levels of IgE antibodies in serum, and test results must be interpreted in the clinical context.

False-positive results for IgE antibodies may occur in patients with markedly elevated serum IgE (>2500 kU/L) due to

nonspecific binding to allergen solid phases.

Clinical Reference

Homburger HA, Hamilton RG: Allergic diseases. In: McPherson RA, Pincus MR, eds. Henry's Clinical Diagnosis and Management by Laboratory Methods. 24th ed. Elsevier; 2022:chap 56

Performance

Method Description

Specific IgE from the patient's serum reacts with the allergen of interest, which is covalently coupled to an ImmunoCAP. After washing away nonspecific IgE, enzyme-labeled anti-IgE antibody is added to form a complex. After incubation, unbound anti-IgE is washed away, and the bound complex incubated with a developing agent. After stopping the reaction, the fluorescence of the eluate is measured. Fluorescence is proportional to the amount of specific IgE present in the patient's sample (ie, the higher the fluorescence value, the more IgE antibody is present).(Package insert: ImmunoCAP System Specific IgE FEIA. Phadia; Rev 02/2024)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

Same day/1 to 3 days

Specimen Retention Time

14 days

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Superior Drive

Fees & Codes

Fees

- Authorized users can sign in to Test Prices for detailed fee information.
- Clients without access to Test Prices can contact Customer Service 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact Customer Service.

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

86003

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
MOLD1	Mold Panel	30183-8

Result ID	Test Result Name	Result LOINC® Value
MOLD1	Mold Panel	30183-8