



Test Definition: ABOPC

Arbovirus Antibody Panel, IgG and IgM, Spinal Fluid

Overview

Useful For

Aiding the diagnosis of arboviral encephalitis due to California (LaCrosse), St. Louis, Eastern equine, or Western equine encephalitis viruses.

Profile Information

| Test Id | Reporting Name | Available Separately | Always Performed |
|---------|-------------------------------------|----------------------|------------------|
| CAVPC | Calif(LaCrosse) Encep Ab Panel, CSF | No | Yes |
| EELPC | East Equine Enceph Ab Panel, CSF | No | Yes |
| STLPC | St. Louis Enceph Ab Panel, CSF | No | Yes |
| WEEPC | West Equine Enceph Ab Panel, CSF | No | Yes |

Testing Algorithm

The following algorithms are available:

[-Meningitis/Encephalitis Panel Algorithm](#)

[-Mosquito-borne Disease Laboratory Testing](#)

Special Instructions

- [Meningitis/Encephalitis Panel Algorithm](#)
- [Mosquito-borne Disease Laboratory Testing](#)

Method Name

Immunofluorescence Assay (IFA)

NY State Available

No

Specimen

Specimen Type

CSF

Ordering Guidance

New York State clients: This test is not available for specimens originating in New York.

Specimen Required

Container/Tube: Sterile vial

Preferred: Vial number 2

Acceptable: Any vial

Specimen Volume: 0.7 mL

Collection Instructions: Submit specimen collected in vial 2, if possible. If not, note which vial from which the aliquot was obtained.

Forms

If not ordering electronically, complete, print, and send [Infectious Disease Serology Test Request](#) (T916) with the specimen.

Specimen Minimum Volume

0.7 mL

Reject Due To

| | |
|-----------------|----|
| Gross hemolysis | OK |
| Gross lipemia | OK |

Specimen Stability Information

| Specimen Type | Temperature | Time | Special Container |
|---------------|--------------------------|---------|-------------------|
| CSF | Refrigerated (preferred) | 14 days | |
| | Frozen | 14 days | |

Clinical & Interpretive**Clinical Information**

California (LaCrosse) Virus:

California (LaCrosse) virus is a member of the Bunyaviridae family, and it is one of the arthropod-borne encephalitides. It is transmitted by various *Aedes* and *Culex* mosquitoes and is found in such intermediate hosts as rabbits, squirrels, chipmunks, and field mice. California meningoencephalitis is usually mild and occurs in late summer. Ninety percent of infections are seen in children and adolescents younger than 15 years, usually from rural areas. The incubation period is estimated to be 7 days, and acute illness lasts 10 days or less in most instances. Typically, the first symptoms are nonspecific, lasting 1 to 3 days, and are followed by the appearance of central nervous system (CNS) signs and symptoms, such as stiff neck, lethargy, and seizures, which usually abate within 1 week. Symptomatic infection is almost never recognized in those older than 18 years. The most important sequela of California virus encephalitis is epilepsy, which occurs in about 10% of children; almost always in patients who have had seizures during the acute illness. An estimated 2% of patients have persistent paresis. Learning disabilities or other objective cognitive deficits have been

reported in a small proportion (2%) of patients. Learning performance and behavior of most recovered patients are not distinguishable from comparison groups in these same areas.

Eastern Equine Encephalitis:

Eastern equine encephalitis (EEE) is within the alphavirus group. It is a low-prevalence cause of human disease in the Eastern and Gulf Coast states. EEE is maintained by a cycle of mosquito/wild bird transmission, peaking in the summer and early fall, when humans may become an adventitious host. The most common clinically apparent manifestation is a mild undifferentiated febrile illness, usually with headache. CNS involvement is demonstrated in only a minority of infected individuals and is more abrupt and more severe than with other arboviruses, with children being more susceptible to severe disease. Fatality rates are approximately 70%.

St Louis Encephalitis:

Areas or outbreaks of St Louis encephalitis since 1933 have involved the western United States, Texas, the Ohio-Mississippi Valley, and Florida. The transmission vector is the mosquito. Peak incidence occurs in summer and early autumn. Disease onset is characterized by generalized malaise, fever, chills, headache, drowsiness, nausea, and sore throat or cough, followed in 1 to 4 days by meningeal and neurologic signs. The severity of illness increases with advancing age; persons older than 60 years have the highest frequency of encephalitis. Symptoms of irritability, sleeplessness, depression, memory loss, and headaches can last up to 3 years.

Western Equine Encephalitis:

The virus that causes Western equine encephalitis (WEE) is widely distributed throughout the United States and Canada; disease occurs almost exclusively in the Western states and Canadian provinces. The relative absence of the disease in the eastern United States probably reflects a paucity of the vector mosquito species, *Culex tarsalis*, and possibly a lower pathogenicity of local virus strains. The disease usually begins suddenly with malaise, fever, and headache, often with nausea and vomiting. Vertigo, photophobia, sore throat, respiratory symptoms, abdominal pain, and myalgia are also common. Over a few days, the headache intensifies; drowsiness and restlessness may merge into a coma in severe cases. In infants and children, the onset may be more abrupt than for adults. WEE should be suspected in any case of febrile CNS disease from an endemic area. Infants are highly susceptible to CNS disease, with about 20% of cases in patients younger than 1 year. There is an excess of male patients with WEE clinical encephalitis, averaging about twice the number of infections detected in female patients. After recovery from the acute disease, patients may require from several months to 2 years to overcome fatigue, headache, and irritability. Infants and children are at a higher risk of permanent brain damage after recovery than adults.

Reference Values**CALIFORNIA VIRUS (La CROSSE) ENCEPHALITIS ANTIBODY**

IgG: <1:1

IgM: <1:1

Reference values apply to all ages.

EASTERN EQUINE ENCEPHALITIS ANTIBODY

IgG: <1:1

IgM: <1:1

Reference values apply to all ages.

ST. LOUIS ENCEPHALITIS ANTIBODY

IgG: <1:1

IgM: <1:1

Reference values apply to all ages.

WESTERN EQUINE ENCEPHALITIS

IgG: <1:1

IgM: <1:1

Reference values apply to all ages.

Interpretation

Detection of organism-specific antibodies in the spinal fluid (CSF) may suggest central nervous system (CNS) infection. However, these results are unable to distinguish between intrathecal antibodies and serum antibodies introduced into the CSF at the time of lumbar puncture or from a breakdown in the blood-brain barrier. The results should be interpreted with other laboratory and clinical data prior to a diagnosis of CNS infection.

Cautions

All results must be correlated with clinical history and other data available to the attending physician.

False-positive results may be caused by breakdown of the blood-brain barrier or by the introduction of blood into the spinal fluid during collection.

Since cross-reactivity with dengue fever virus does occur with St Louis encephalitis antigens and, therefore, cannot be differentiated further, the specific virus responsible for positive results may be deduced by the travel history of the patient, along with available medical and epidemiological data, unless the virus can be isolated.

Eastern and Western equine encephalitis viruses show some cross-reactivity; however, antibody response to the infecting virus is typically at least 8-fold higher.

Clinical Reference

Piantadosi A, Kanjilal S. Diagnostic approach for arboviral infections in the United States. *J Clin Microbiol.* 2020;58(12):e01926-19. doi:10.1128/JCM.01926-19

Performance**Method Description**

The indirect immunofluorescent antibody (IFA) assay is a 2-stage sandwich procedure. In the first stage, the patient spinal fluid is diluted in Pretreatment Diluent for IgM and phosphate buffered saline (PBS) for IgG, added to appropriate slide wells in contact with the substrate, and incubated. Following incubation, the slide is washed in PBS, which removes unbound CSF antibodies. In the second stage, each antigen well is overlaid with fluorescein-labeled antibody to IgM and IgG. The slide is incubated allowing antigen-antibody complexes to react with the fluorescein-labeled anti-IgM and anti-IgG. After the slide is washed, dried, and mounted, it is examined using fluorescence microscopy. Positive reactions appear as cells exhibiting bright apple-green cytoplasmic fluorescence against a background of red negative control cells.

Semi-quantitative endpoint titers are obtained by testing serial dilutions of positive specimens.(Package inserts: Arbovirus IFA IgM and Arbovirus IFA IgG Instructions for Use. Focus Diagnostics; Rev 03, 02/17/2023)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

Same day/1 to 4 days

Specimen Retention Time

2 weeks

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Superior Drive

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

86651 x 2-California Virus (La Crosse) Encephalitis Antibody Panel, IgG and IgM, Spinal Fluid

86652 x 2-Eastern Equine Encephalitis Antibody Panel, IgG and IgM, Spinal Fluid

86653 x 2-St. Louis Encephalitis Antibody Panel, IgG and IgM, Spinal Fluid

86654 x 2-Western Equine Encephalitis Antibody Panel, IgG and IgM, Spinal Fluid

LOINC® Information

| Test ID | Test Order Name | Order LOINC® Value |
|---------|-------------------------------------|--------------------|
| ABOPC | Arbovirus Ab Panel IgG and IgM, CSF | 49094-6 |

| Result ID | Test Result Name | Result LOINC® Value |
|-----------|-----------------------------------|---------------------|
| 26365 | Calif(LaCrosse) Encep Ab, IgG,CSF | 9539-8 |
| 26366 | Calif(LaCrosse) Encep Ab, IgM,CSF | 9540-6 |

Test Definition: ABOPC

Arbovirus Antibody Panel, IgG and IgM, Spinal
Fluid

| | | |
|-------|---------------------------------|---------|
| 26369 | East Equine Enceph Ab, IgG, CSF | 10897-7 |
| 26370 | East Equine Enceph Ab, IgM, CSF | 10899-3 |
| 26367 | St. Louis Enceph Ab, IgG, CSF | 21509-5 |
| 26368 | St. Louis Enceph Ab, IgM, CSF | 21510-3 |
| 26371 | West Equine Enceph Ab, IgG, CSF | 9315-3 |
| 26372 | West Equine Enceph Ab, IgM, CSF | 9316-1 |