

## Overview

### Useful For

Monitoring chlordiazepoxide therapy

Assessing toxicity

### Method Name

Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)

### NY State Available

Yes

## Specimen

### Specimen Type

Serum Red

### Shipping Instructions

Ship specimen in amber vial to protect from light.

### Specimen Required

**Supplies:** Amber Frosted Tube, 5 mL (T915)

**Collection Container/Tube:** Red top (serum get/SST are **not acceptable**)

**Submission Container/Tube:** Amber vial

**Specimen Volume:** 1 mL

**Collection Instructions:** Within 2 hours of collection, centrifuge and aliquot serum into an amber vial.

### Forms

If not ordering electronically, complete, print, and send 1 of the following forms with the specimen:

-[Neurology Specialty Testing Client Test Request](#) (T732)

-[Therapeutics Test Request](#) (T831)

### Specimen Minimum Volume

0.3 mL

### Reject Due To

Gross hemolysis	OK
Gross lipemia	OK
Gross icterus	OK

**Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Serum Red	Refrigerated (preferred)	28 days	LIGHT PROTECTED
	Ambient	28 days	LIGHT PROTECTED
	Frozen	28 days	LIGHT PROTECTED

**Clinical & Interpretive****Clinical Information**

Chlordiazepoxide (Librium) is a benzodiazepine widely used in the treatment of anxiety and alcohol withdrawal symptoms, as well as a premedication for anesthesia. The mechanism of action of all benzodiazepines remains unclear. However, it is known that benzodiazepines facilitate gamma-amino butyric acid (GABA)-mediated neurotransmission in the brain. Benzodiazepines most likely facilitate the inhibitory presynaptic or postsynaptic reactions of GABA.

Chlordiazepoxide is metabolized to long-acting metabolites in the liver to the active metabolites nordiazepam (desmethyldiazepam) and oxazepam, and the clearance of the drug is reduced considerably in older adults and in patients with hepatic disease.

Therapeutic assessment should include measurement of both the parent drug (chlordiazepoxide) and the primary active metabolite (nordiazepam), with a proposed combined chlordiazepoxide and nordiazepam reference interval of 700 to 1000 ng/mL. However, therapeutic and toxic concentrations can vary greatly and should be interpreted in the full clinical context of the patient. Since chlordiazepoxide has a wide therapeutic index and toxicity is dose-dependent, routine drug monitoring is not indicated in all patients.

**Reference Values**

Therapeutic concentration:

Chlordiazepoxide: 400-3,000 ng/mL

Cutoff concentrations by liquid chromatography tandem mass spectrometry:

Chlordiazepoxide: 10 ng/mL

Nordiazepam: 10 ng/mL

Oxazepam: 10 ng/mL

**Interpretation**

Combined concentrations of chlordiazepoxide and nordiazepam above 5000 ng/mL have been associated with toxicity.

**Cautions**

The specimen must be protected from light.

**Clinical Reference**

1. Langman LJ, Bechtel LK, Holstege CP. Clinical toxicology. In: Rifai N, Chiu RWK, Young I, Burnham CAD, Wittwer CT, eds. Tietz Textbook of Laboratory Medicine. 7th ed. Elsevier; 2023:chap 43

2. Burtis CA, Ashwood ER, Bruns DE, eds. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics. WB Saunders Company; 2011:1109-1188
3. Hiemke C, Bergemann N, Clement HW, et al. Consensus guidelines for therapeutic drug monitoring in neuropsychopharmacology: Update 2017. *Pharmacopsychiatry*. 2018;51(1-02):9-62
4. Adeli K, Higgins V, Bohn MK. Reference information for the clinical laboratory. In: In: Rifai N, Chiu RWK, Young I, Burnham CAD, Wittwer CT, eds. Tietz Textbook of Laboratory Medicine. 7th ed. Elsevier; 2023:1390-1470

## Performance

### Method Description

The internal standard mixture containing chlordiazepoxide-d5, diazepam-d5, nordiazepam-d5, oxazepam-d5, and temazepam-d5 is added to serum samples, mixed, and centrifuged. The supernatant is diluted and injected on a liquid chromatography tandem mass spectrometer.(Unpublished Mayo method)

### PDF Report

No

### Day(s) Performed

Wednesday

### Report Available

2 to 7 days

### Specimen Retention Time

14 days

### Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Superior Drive

## Fees & Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

### Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

### CPT Code Information

80299

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
CDP	Chlordiazepoxide and metabolites, S	33060-5

Result ID	Test Result Name	Result LOINC® Value
8610	Chlordiazepoxide	3457-9
37321	Nordiazepam	3537-8
622870	Oxazepam	3886-9