

**Test Definition: FCELG** 

Celery IgG

Reporting Title: Celery IgG

Performing Location: VIRACOR-IBT LABORATORIES, INTERFACE

## **Specimen Requirements:**

Draw blood in a plain red-top tube(s), serum gel tube(s) is acceptable. Spin down and send 0.5 mL of serum in a plastic vial.

Specimen Type	Temperature	Time
Serum	Refrigerated (preferred)	28 days
	Frozen	365 days
	Ambient	7 days

## **Result Codes:**

Result ID	Reporting Name	Туре	Unit	LOINC®
FCELG	Celery IgG	Alphanumeric		In Process

## **CPT Code Information:**

86001

## Reference Values:

<2 mcg/mL

The reference range listed on the report is the lower limit of quantitation for the assay. The clinical utility of food-specific IgG tests has not been established. These tests can be used in special clinical situations to select foods for evaluation by diet elimination and challenge in patients who have food-related complaints. It should be recognized that the presence of food-specific IgG alone cannot be taken as evidence of food allergy and only indicates immunologic sensitization by the food allergen in question. This test should only be ordered by physicians who recognize the limitations of the test.

Test Performed by: Viracor-IBT Laboratories

1001 NW Technology Dr. Lee's Summit, MO 64086