

Spinocerebellar Ataxia Repeat Expansion Panel, Varies

Test ID: SCAP

Explanation: On the effective date extracted DNA will be acceptable for testing. Formatting of acceptable specimen types will also be standardized. Reflex testing will be updated to include MATCC, Maternal Cell Contamination testing for cord blood specimens.

| Current Algorithm |
|-------------------|
| None |

| New Algorithm |
|--|
| For cord blood specimens that have an accompanying maternal blood specimen, maternal cell contamination studies will be performed at an additional charge. |

| Current Reflex Tests |
|----------------------|
| None |

| New Reflex Tests | | | |
|------------------|--------------------------------|----------------------|------------------|
| Test ID | Reporting Name | Available Separately | Always Performed |
| MATCC | Maternal Cell Contamination, B | Yes | No |

| Current Specimen Information | |
|------------------------------|--------------------------|
| Specimen Type | Temperature |
| Varies | Refrigerated (preferred) |
| | Ambient |
| | Frozen |

| New Specimen Information | |
|--------------------------|-------------|
| Specimen Type | Temperature |
| Varies | Varies |

| Current Specimen Required | New Specimen Required |
|---|--|
| <p>Specimen Type: Whole blood</p> <p>Patient Preparation: A previous bone marrow transplant from an allogenic donor will interfere with testing. Call 800-533-1710 for instructions for testing patients who have received a bone marrow transplant.</p> <p>Container/Tube:</p> <p>Preferred: Lavender top (EDTA) or yellow top (ACD)</p> <p>Acceptable: Any anticoagulant</p> <p>Specimen Volume: 3 mL</p> <p>Collection Instructions:</p> <ol style="list-style-type: none"> 1. Invert several times to mix blood. 2. Send whole blood specimen in original tube. Do not aliquot. | <p>Patient Preparation: A previous hematopoietic stem cell transplant from an allogenic donor will interfere with testing. For information about testing patients who have received a hematopoietic stem cell transplant, call 800-533-1710.</p> <p>Submit only 1 of the following specimens:</p> <p>Specimen Type: Whole blood</p> <p>Container/Tube:</p> <p>Preferred: Lavender top (EDTA) or yellow top (ACD)</p> <p>Specimen Volume: 3 mL</p> <p>Collection Instructions:</p> <ol style="list-style-type: none"> 1. Invert several times to mix blood. 2. Send whole blood specimen in original tube. Do not aliquot. 3. Whole blood collected postnatal from an umbilical cord is also acceptable. See Additional Information <p>Specimen Stability Information: Ambient (preferred) 4 days/Refrigerated 4 days/Frozen 4 days</p> <p>Additional Information:</p> <ol style="list-style-type: none"> 1. Specimens are preferred to be received within 4 days of collection. Extraction will be attempted for specimens received after 4 days, and DNA yield will be evaluated to determine if testing may proceed. 2. To ensure minimum volume and concentration of DNA are met, the requested volume must be submitted. Testing may be canceled if DNA requirements are inadequate. 3. For postnatal umbilical cord whole blood specimens, maternal cell contamination studies are recommended to ensure test results reflect that of the patient tested. A maternal blood specimen is required to complete maternal cell contamination studies. Order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on both the cord blood and maternal blood specimens under separate order numbers. <p>Specimen Type: Extracted DNA</p> <p>Container/Tube:</p> <p>Preferred: Screw Cap Micro Tube, 2mL with skirted conical base</p> <p>Acceptable: Matrix tube, 1 mL</p> <p>Collection Instructions:</p> <ol style="list-style-type: none"> 1. The preferred volume is at least 100 mcL at a concentration of 75 ng/mcL. 2. Include concentration and volume on tube. <p>Specimen Stability Information: Frozen (preferred) 1 year/Ambient/Refrigerated</p> <p>Additional Information: DNA must be extracted in a CLIA-certified laboratory or equivalent and must be extracted from a specimen type listed as acceptable for this test (including applicable anticoagulants). Our laboratory has experience with Chemagic, Puregene, Autopure, MagnaPure, and EZ1 extraction platforms and cannot guarantee that all extraction methods are compatible with this test. If testing fails, one repeat will be attempted, and if unsuccessful, the test will be reported</p> |

as failed and a charge will be applied. If applicable, specific gene regions that were unable to be interrogated due to DNA quality will be noted in the report.

Current Specimen Retention Time

Whole Blood: 2 weeks (if available), Extracted DNA: 3 months

New Specimen Retention Time

Whole blood: 28 days (if available); Extracted DNA: 3 months

Current Days Performed

Monday, Wednesday

New Days Performed

Varies

Current Specimen Retention Time

Whole blood: 2 weeks (if available) Extracted DNA: 3 months

New Specimen Retention Time

Whole blood: 28 days (if available); Extracted DNA: 3 months

Questions

Contact Melissa Tricker-Klar, Laboratory Resource Coordinator at 800-533-1710.

© Mayo Foundation for Medical Education and Research. All rights reserved.