

Ehlers-Danlos Syndrome Gene Panel, Varies

Test ID: EDSGG

Explanation: On the effective date extracted DNA will be acceptable for testing. Formatting of acceptable specimen types will also be standardized. Reflex testing will be updated to include MATCC, Maternal Cell Contamination testing for cord blood specimens.

Current Ordering Guidance
Customization of this panel and single gene analysis for any gene present on this panel are available. For more information see CGPH/ Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies. Targeted testing for familial variants (also called site-specific or known mutations testing) is available for the genes on this panel. See FMTT / Familial Variant, Targeted Testing, Varies. To obtain more information about this testing option, call 800-533-1710.

New Ordering Guidance
Customization of this panel and single gene analysis for any gene present on this panel are available. For more information see CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies. To modify this panel via CGPH, use the Cardiovascular/Connective Tissue/Dyslipidemia/Cerebrovascular/Primary Ciliary Dyskinesia disease state for step 1 on the Custom Gene Ordering Tool . Targeted testing for familial variants (also called site-specific or known mutations testing) is available for the genes on this panel. See FMTT / Familial Variant, Targeted Testing, Varies. To obtain more information about this testing option, call 800-533-1710.

Current Reflex Tests
None

New Reflex Tests			
Test ID	Reporting Name	Available Separately	Always Performed
MATCC	Maternal Cell Contamination, B	Yes	No

Current Specimen Required	New Specimen Required
<p>Patient Preparation: A previous bone marrow transplant from an allogenic donor will interfere with testing. For instructions for testing patients who have received a bone marrow transplant, call 800-533-1710</p> <p>Specimen Type: Whole blood</p> <p>Container/Tube:</p> <p>Preferred: Lavender top (EDTA) or yellow top (ACD)</p> <p>Acceptable: Any anticoagulant</p> <p>Specimen Volume: 3 mL</p> <p>Collection Instructions:</p> <ol style="list-style-type: none"> 1. Invert several times to mix blood. 2. Send whole blood specimen in original tube. Do not aliquot. <p>Specimen Stability Information: Ambient (preferred)/Refrigerated</p> <p>Specimen Type: Saliva</p> <p>Patient Preparation: Patient should not eat, drink, smoke, or chew gum 30 minutes prior to collection.</p> <p>Supplies:</p> <p>DNA Saliva Kit High Yield (T1007) Saliva Swab Collection Kit (T786)</p> <p>Container/Tube:</p> <p>Preferred: High-yield DNA saliva kit</p> <p>Acceptable: Saliva swab</p> <p>Specimen Volume: 1 Tube if using T1007 or 2 swabs if using T786</p> <p>Collection Instructions: Collect and send specimen per kit instructions.</p> <p>Specimen Stability Information: Ambient (preferred) 30 days/Refrigerated 30 days</p> <p>Additional Information: Saliva specimens are acceptable but not recommended. Due to lower quantity/quality of DNA yielded from saliva, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.</p>	<p>Patient Preparation: A previous hematopoietic stem cell transplant from an allogenic donor will interfere with testing. For information about testing patients who have received a hematopoietic stem cell transplant, call 800-533-1710.</p> <p>Submit only 1 of the following specimens:</p> <p>Specimen Type: Whole blood</p> <p>Container/Tube: Lavender top (EDTA) or yellow top (ACD)</p> <p>Specimen Volume: 3 mL</p> <p>Collection Instructions:</p> <ol style="list-style-type: none"> 1. Invert several times to mix blood. 2. Send whole blood specimen in original tube. Do not aliquot. 3. Whole blood collected postnatal from an umbilical cord is also acceptable. See Additional Information <p>Specimen Stability Information: Ambient (preferred) 4 days/Refrigerated 4 days/Frozen 4 days</p> <p>Additional Information:</p> <ol style="list-style-type: none"> 1. Specimens are preferred to be received within 4 days of collection. Extraction will be attempted for specimens received after 4 days, and DNA yield will be evaluated to determine if testing may proceed. 2. To ensure minimum volume and concentration of DNA are met, the requested volume must be submitted. Testing may be canceled if DNA requirements are inadequate. 3. For postnatal umbilical cord whole blood specimens, maternal cell contamination studies are recommended to ensure test results reflect that of the patient tested. A maternal blood specimen is required to complete maternal cell contamination studies. Order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on both the cord blood and maternal blood specimens under separate order numbers. <p>Specimen Type: Saliva</p> <p>Patient Preparation: Patient should not eat, drink, smoke, or chew gum 30 minutes prior to collection.</p> <p>Supplies:</p> <p>DNA Saliva Kit High Yield (T1007) Saliva Swab Collection Kit (T786)</p> <p>Container/Tube:</p> <p>Preferred: High-yield DNA saliva kit</p> <p>Acceptable: Saliva swab</p> <p>Specimen Volume: 1 Tube if using T1007 or 2 swabs if using T786</p> <p>Collection Instructions: Collect and send specimen per kit instructions.</p> <p>Specimen Stability Information: Ambient (preferred) 30 days/Refrigerated 30 days</p> <p>Additional Information: Saliva specimens are acceptable but not recommended. Due to lower quantity/quality of DNA yielded from saliva, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific</p>

gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.

Specimen Type: Extracted DNA

Container/Tube:

Preferred: Screw Cap Micro Tube, 2mL with skirted conical base

Acceptable: Matrix tube, 1 mL

Collection Instructions:

1. The preferred volume is at least 100 mcL at a concentration of 75 ng/mcL.

2. Include concentration and volume on tube.

Specimen Stability Information: Frozen (preferred) 1 year/Ambient/Refrigerated

Additional Information: DNA must be extracted in a CLIA-certified laboratory or equivalent and must be extracted from a specimen type listed as acceptable for this test (including applicable anticoagulants). Our laboratory has experience with Chemagic, Puregene, Autopure, MagnaPure, and EZ1 extraction platforms and cannot guarantee that all extraction methods are compatible with this test. If testing fails, one repeat will be attempted, and if unsuccessful, the test will be reported as failed and a charge will be applied. If applicable, specific gene regions that were unable to be interrogated due to DNA quality will be noted in the report.

Current Report Available

28 to 42 days

New Report Available

21 to 28 days

Current Specimen Retention Time

Whole blood: 2 weeks (if available); Extracted DNA: 3 months; Saliva: 1 month

New Specimen Retention Time

Whole blood: 28 days (if available); Saliva: 30 days (if available); Extracted DNA: 3 months

Questions

Contact Melissa Tricker-Klar, Laboratory Resource Coordinator at 800-533-1710.